# CONTENTS


| | | PAGE |
|---|---|---|
| Post-harvest dormancy in wheat (*Triticum aestivum*) collections from Nepal hills | Jag Shoran, J. P. Saini and H. C. Joshi | 1 |
| Genetic diversity in relation to geographic diversity in maize (*Zea mays*) | S. K. Prasad, T. P. Singh and M. N. Haque | 4 |
| Combining ability by line × tester analysis in castor (*Ricinus communis*) | U. G. Fatteh, C. J. Dangaria, K. L. Dobariya and V. J. Patel | 7 |
| Combining-ability analysis for quantitative traits of rocket-salad (*Eruca vesicaria* subsp. *sativa*) grown on normal and alkaline soils | D. Kumar, I. S. Yadav and B. S. Gupta | 11 |
| Bases of resistance in okra (*Abelmoschus esculentus*) to *Amrasca biguttula biguttula* | Ram Singh | 15 |
| Effect of intercropping of legumes in sorghum (*Sorghum bicolor*) at different levels of nitrogen under rainfed condition on sorghum equivalent and monetary return | M. L. Gupta and R. M. Singh | 20 |
| Post-monsoon irrigation of pigeonpea (*Cajanus cajan*) in intercropping system in black soils of Tawa-command area | K. P. Tiwari, J. P. Dixit and R. N. Saran | 26 |
| Interrelations of evapotranspiration from wheat (*Triticum aestivum*) and free water surface and open-pan evaporation in north India | V. K. Arora, S. S. Prihar and P. R. Gajri | 31 |
| Interrelationship between rainfall and rice green leafhopper population in West Bengal | Sujata Mukhopadhyay, P. S. Nath, T. K. Sarkar, S. Sarkar and S. Mukhopadhyay | 34 |
| Termination of larval diapause and its importance in forecasting incidence of soybean girdler, *Obereopsis brevis* (Coleoptera : Lamiidae) | N. Dutt and P. K. Pal | 38 |

| | | PAGE |
|---|---|---|
| Environmental index for determining the adaptation of crops | R. K. Mahajan and A. S. R. Prasad | 42 |
| Design, development and evaluation of two-row tractor-mounted ridge-planter for winter maize (*Zea mays*) | Surendra Singh, V. K. Sharma, P. K. Gupta and A. M. Chauhan | 45 |
| Variability and character association in blackgram (*Vigna mungo*) | K. B. Wanjari | 48 |
| Correlation and discriminant function analysis in Indian mustard (*Brassica juncea* subsp. *juncea*) | Narendra Kumar, J. K. Bisht and M. C. Joshi | 51 |
| Effect of pollen quantity on pod and seed setting in *Brassica* species | R. K. Katiyar | 52 |
| Variety × season interaction in sunflower (*Helianthus annuus*) | S. K. Chaudhary and I. J. Anand | 55 |
| Correlation and path analysis in cowpea (*Vigna unguiculata* subsp. *cylindrica*) | P. C. Tyagi and K. D. Koranne | 57 |
| Genetic variation in Persian clover (*Trifolium resupinatum*) | R. K. Gumber and M. S. Sohoo | 58 |
| Variability and correlation in palmarosa (*Cymbopogon martinii*) | M. S. Punia, P. K. Verma and G. D. Sharma | 59 |
| Genotype × environment interaction for cured-leaf yield and earliness in cigar-wrapper tobacco (*Nicotiana tabacum*) | V. K. Dobhal | 62 |
| Genetics of quantitative characters in tomato (*Lycopersicon lycopersicum*) | N. D. Das, S. S. Ghosh and T. K. Chattopadhyay | 64 |
| Land-equivalent ratio in evaluation of mixed-cropping, intercropping and sole-cropping systems | Syed Moinbasha and R. A. Singh | 65 |
| Response of summer groundnut (*Arachis hypogaea*) cultivars to sources and levels of phosphorus | G. L. Maliwal, N. K. Tank and D. N. Kunadia | 67 |
| Calcium content of potato (*Solanum tuberosum*) plant as affected by potassium application | U. C. Sharma and B. R. Arora | 69 |
| Effect of suppression of flowering on improvement of growth, yield and juice quality in sugarcane | Sudama Singh, K. Mohan Naidu and D. N. Tyagi | 71 |
| Prediction of dry-matter accumulation through non-destructive methods in papaya (*Carica papaya*) | K. Balakrishnan, K. M. Sundaram, N. Natarajaratnam and C. Rajendran | 74 |

| | | PAGE |
|---|---|---|
| Culture of *Beauveria bassiana* and its pathogenicity to insect pests of jute (*Corchorus capsularis* and *C. olitorius*) and mesta (*Hibiscus cannabinus* and *H. sabdariffa*) | N. C. Pandit and D. Som | 75 |
| Post-treatment temperature effect on toxicity of some acaricides to *Tetranychus neocaledonicus* on okra (*Abelmoschus esculentus*) | M. P. Singh, D. D. Sharma and S. F. Hameed | 77 |
| Field evaluation of various chemicals for the control of leopard moth, *Zeuzera multistrigata*, on cherry (*Prunus avium*) | S. P. Bhardwaj | 79 |
| Efficacy of plastic-house in propagation of guava (*Psidium guajava*) through hardwood cuttings | K. Malla Reddy and R. N. Singh | 81 |
| Development of an animal-drawn, inclined-roller crust breaker | N. K. Awadhwal | 87 |
| Evaluation of leaf-area index in greengram (*Vigna radiata*) | R. Nanda and A. D. Saini | 91 |
| Contribution of area, thickness and conductance of leaf in biomass production in upland cotton (*Gossypium hirsutum*) | S. N. Bhardwaj, A. D. Saini, Munshi Singh and K. P. Singh | 100 |
| Genetics of grain yield and drought index in pearlmillet (*Pennisetum typhoides*) under moisture stress | Prem Sagar and R. L. Kapoor | 104 |
| Outcrossing for improvement in yield and quality of upland cotton (*Gossypium hirsutum*) in north India | Munshi Singh, V. P. Singh, Kamal Paul and C. B. Lal | 110 |
| Genetic analysis of some quantitative traits in hookah and chewing tobacco (*Nicotiana rustica*) | V. K. Dobhal | 114 |
| Sources of resistance to hill bunt in wheat (*Triticum aestivum*) | S. R. Singh and V. L. Chopra | 119 |
| Resistance of groundnut (*Arachis hypogaea*) to aflatoxin | Kiran Kalia, H. M. Desai and M. K. Chakraborty | 121 |
| Inheritance of disease resistance in blackgram (*Vigna mungo*) to mungbean yellow mosaic virus | R. P. Kaushal and B. M. Singh | 123 |

| | | PAGE |
|---|---|---|
| Analysis of light-trap catches of *Nephotettix* species (Hemiptera : Cicadellidae) in West Bengal | John Bowden, Sujata Mukhopadhyay, P. S. Nath, T. K. Sarkar, S. Sarkar and S. Mukhopadhyay | 125 |
| Potato tuber moth, *Phthorimaea operculella* (Zeller), in North-Eastern Hills region and a simple method for its control | Lakshman Lal | 130 |
| Rainfall probability analysis of 3 stations in Andhra Pradesh for crop planning | A. Yogeswara Rao, K. Krishna Rao and B. V. Ramana Rao | 133 |
| Effect of planting date and hormone on onion (*Allium cepa*) yield | S. P. S. Tomar, T. P. Singh and G. S. Tomar | 136 |
| Stability analysis for phenological traits in chickpea (*Cicer arietinum*) | K. P. Singh, C. S. Tyagi, B. D. Chaudhary and V. P. Singh | 139 |
| Genetic variability for peg strength and related characters in groundnut (*Arachis hypogaea*) | K. Johnson George, S. P. Tiwari and P. S. Reddy | 141 |
| A complete fertility-restorer line in upland cotton (*Gossypium hirsutum*) | Sunil Seth, D. K. Jain and M. S. Kairon | 144 |
| Vesicular-arbuscular mycorrhizal symbiosis in coconut (*Cocos nucifera*) in relation to the root (wilt) disease and intercropping or mixed cropping | George V. Thomas | 145 |
| Fungicidal control of anthracnose and fruit-rot (*Colletotrichum capsici*) of chilli (*Capsicum annuum*) | D. Mishra | 147 |
| Biology of an ichneumonid *Hyposoter didymator* on *Spodoptera litura* | Pradyumn Kumar, S. P. Singh, S. K. Jalali and Chandish R. Ballal | 149 |
| Effect of edible oils in protection of greengram (*Vigna radiata*) seed from pulse beetles (*Callosobruchus chinensis* and *C. maculatus*) | R. B. Doharey, R. N. Katiyar and K. M. Singh | 151 |
| Heptachlor uptake by potato (*Solanum tuberosum*), sweet-potato (*Ipomoea batatas*) and onion (*Allium cepa*) from treated soils | V. J. Shivankar and V. S. Kavadia | 155 |
| Combining-ability analysis of reaction of sorghum (*Sorghum bicolor*) genotype to *Striga asiatica* | N. B. Barche, A. R. Dabholkar, R. C. Mishra and G. S. Lal | 161 |

| | | PAGE |
|---|---|---|
| Genetic diversity in relation to plant type in early pigeonpea (*Cajanus cajan*) | S. P. Singh, J. N. Govil, Hayat Ram and R. B. Mehra | 165 |
| Genetic analysis of nitrogen-fixation traits in greengram (*Vigna radiata*) | B. D. Singh and B. K. Murty | 171 |
| Genetic analysis of quantitative characters in a cross of Indian mustard (*Brassica juncea* subsp. *juncea*) | J. N. Sachan and Basudeo Singh | 176 |
| Resistance of sesame (*Sesamum indicum*) varieties to gallfly (*Asphondylia sesami*) and capsule-borer (*Antigastra catalaunalis*) | K. Bhagavan Dora and T. Kamala | 180 |
| Effect of gamma-rays on heterosis in cotton (*Gossypium arboreum* and *G. hirsutum*) hybrids | S. S. Duhoon and P. V. Verma | 184 |
| Potential for seed oil in germplasm of tree cotton (*Gossypium arboreum*) and Egyptian cotton (*G. arboreum* var. *nadam*) | Phundan Singh | 188 |
| Influence of nitrogen, zinc and their interaction on the yield and nutrient uptake of 'IR 20' rice (*Oryza sativa*) in different seasons | M. Abdul Salam and S. Subramanian | 190 |
| Production potential and economic profitability of different cropping patterns including cereals, pulses and oilseeds | H. C. Thakur and N. N. Sharma | 194 |
| Improvement of monetary return of coriander (*Coriandrum sativum*) through safflower (*Carthamus tinctorius*) intercropping in dryland | S. M. Nikam, M. S. Mahajan and A. B. Deokar | 202 |
| Effect of seed bacterization with *Azotobacter chroococcum* on yield of upland cotton (*Gossypium hirsutum*) | S. T. Shende, Munshi Singh and V. P. Singh | 206 |
| Stage-dependent differential effect of saline water irrigation on grain yield and chemical composition of pearl-millet (*Pennisetum typhoides*) | K. S. Gill | 210 |
| Stomatal responses of cocoa (*Theobroma cacao*) to climatic factors | D. Balasimha and V. Rajagopal | 213 |
| Assessment of incidence and crop loss due to pod-borers of pigeonpea (*Cajanus cajan*) of different maturity groups | C. P. Yadava, S. S. Lal and J. N. Sachan | 216 |

| | | PAGE |
|---|---|---|
| Tolerance of wheats (*Triticum* species) to metoxuron | J. K. Misra and R. K. S. Rathore | 219 |
| Maximization of yield of rainfed pearl-millet (*Pennisetum typhoides*) | R. C. Gautam and S. K. Kaushik | 223 |
| Effect of nitrogen, boron and molybdenum on yield of root and leaf protein of radish (*Raphanus sativus*) | D. K. Bagchi, A. Banerjee and B. Bhattacharya | 224 |
| Interactions between *Azospirillum* species and wheat (*Triticum aestivum*) varieties | Indu Bala and B. S. Kundu | 227 |
| Effect of zinc and copper and *Rhizobium* inoculation on nodulation and yield of greengram (*Vigna radiata*) | Jagdish Prasad and H. Ram | 230 |
| Field reaction of apple (*Malus pumila*) germplasm to powdery mildew (*Podosphaera leucotricha*) | K. D. Verma and G. K. Gupta | 233 |
| Interaction of *Meloidogyne incognita* and *Rhizoctonia bataticola* in white jute (*Corchorus capsularis*) | C. Mishra, D. Som and B. Singh | 234 |
| Exit behaviour of pink bollworm (*Pectinophora gossypiella*) larvae | J. P. Singh, B. P. S. Lather and B. R. Mor | 236 |
| Diflubenzuron and neem (*Azadirachta indica*) oil in control of *Heliothis armigera* infesting chickpea (*Cicer arietinum*) | S. N. Sinha and K. N. Mehrotra | 238 |
| Contact toxicity of penfluron on *Pericallia ricini* larvae | B. B. L. Srivastava and M. Maqsood Khan | 240 |
| Aldicarb residues in coriander (*Coriandrum sativum*) grown on treated soil | P. C. Jain, C. P. S. Yadav and B. L. Pareek | 241 |
| Dispersal of milky-disease organism *Bacillus popilliae* var. *holotrichiae* of white grub (*Holotrichia consanguinea*) through birds | R. V. Vyas, B. M. Parasharya and D. N. Yadav | 243 |
| Linkage relationship for seedling resistance to races of stem-rust (*Puccinia graminis* f. sp. *tritici*) in tetraploid wheats (*T. durum* and *T. carthlicum*) | V. M. Raut, V. P. Patil and G. B. Deodikar | 247 |
| Combining-ability analysis for oil content and iodine value in linseed (*Linum usitatissimum*) | K. S. Karmuka, Tashi Dawa and G. S. Sethi | 252 |

| | | PAGE |
|---|---|---|
| Evolution of upland cotton (*Gossypium hirsutum*) lines with multi-racial resistance to bacterial blight caused by *Xanthomonas campestris* pv. *malvacearum* | V. V. Singh, M. K. Meshram, Sheo Raj and S. S. Narayanan | 255 |
| Intercropping of maize (*Zea mays*) with short-duration pigeonpea (*Cajanus cajan*) and groundnut (*Arachis hypogaea*) | H. C. Thakur and N. N. Sharma | 259 |
| Dry-matter accumulation and yield of winter French-bean (*Phaseolus vulgaris*) as influenced by genotype, nitrogen level and plant population | Masood Ali and A. Tripathi | 263 |
| Feasibility of minimum tillage and cultural practices in potato (*Solanum tuberosum*) cultivation | A. K. Bhattacharjee and V. S. Kushwah | 267 |
| Effect of different moisture regimes and soil test-crop response of nitrogen on yield of rice (*Oryza sativa*) | T. Yellamanda Reddy and R. Kulandaivelu | 274 |
| Forage yield, chemical composition and reclaiming effects of Karnal grass (*Diplachne fusca*) grown with trees in alkali soil | S. S. Grewal and I. P. Abrol | 278 |
| Temporal and genotypic variations in the storability of rice (*Oryza sativa*) under the ambient conditions of Kerala | B. Mohan Kumar and V. P. Sukumara Dev | 283 |
| Assessment of avoidable loss in cotton (*Gossypium hirsutum* and *G. arboreum*) due to sucking pests and bollworms | Ashok K. Dhawan, A. S. Sidhu and G. S. Simwat | 290 |
| Phenology and chemical control of aphid *Myzus persicae* on henbane (*Hyoscyamus niger* and *H. muticus*) | D. Singh, A. K. Tripathi and S. M. Rao | 292 |
| Evaluation of weather-proof baits for the control of field rodents in rainy season | N. Ahmad, C. Sheikher and S. S. Guraya | 297 |
| Transgressive traits in interspecific crosses of wheat (*Triticum* species) | S. N. Nayak | 299 |
| Combining-ability analysis in durum wheat (*Triticum durum*) | R. K. Chowdhury and S. S. Dhanda | 301 |

| | | PAGE |
|---|---|---|
| Interrelationships of quantitative characters with seed yield in $F_2$, $F_3$, $F_4$ and $F_5$ generations of soybean (*Glycine max*) | Pushpendra and Hari Har Ram | 304 |
| Inheritance and association of some quantitative traits in a diallel set of upland cotton (*Gossypium hirsutum*) | H. R. Garg and H. S. Kalsy | 306 |
| Variability and correlations for boll attributes in upland cotton (*Gossypium hirsutum*) germplasm | V. V. Singh | 309 |
| Albinism in segregating seedlings of potato (*Solanum tuberosum*) | K. R. Dhiman and B. L. Barua | 310 |
| Radiation-induced stout fruit mutants of chilli (*Capsicum annuum*) | N. Lakshmi and N. B. Rao | 311 |
| Effect of salinity and gamma-ray treatment on alpha-amylase activity in 'HD 2009' bread wheat (*Triticum aestivum*) | V. Kumar and D. Kumar | 313 |
| Pre-planting soaking treatment of seed tubers for increasing yield of potato (*Solanum tuberosum*) | R. C. Sharma | 315 |
| Morphology and yield of chilli (*Capsicum annuum*) in relation to distance from lemon-scented eucalyptus (*Eucalyptus citriodora*) stands | A. B. I. Igboanugo | 317 |
| Effect of growth-regulators on sugars in sugarcane juice | Ashok Kumar and H. P. Pande | 319 |
| Estimation of the area of detached or intact leaves of papaya (*Carica papaya*) | I. O. O. Aiyelaagbe and M. O. A. Fawusi | 322 |
| Seed-borne infection of bacterial blight (*Xanthomonas campestris* pv. *malvacearum*) in cotton (*Gossypium* species) | N. K. Taneja and Sheo Raj | 323 |
| Occurrence of stem-end rot of Kinnow mandarin (*Citrus reticulata*) and its control through post-harvest treatment with fungicides | B. A. Ullasa and R. D. Rawal | 324 |
| Effects of three synthetic pyrethroids and carbaryl on the incidence of bollworms and yield of cotton (*Gossypium hirsutum*) | J. P. Sharma and R. A. Agarwal | 326 |

| | | PAGE |
|---|---|---|
| Laboratory evaluation of anticoagulant wax block to control Indian field-mouse (*Mus booduga*) | M. Balasubramanyam, G. P. Maddaiah and K. R. Purushotham | 330 |
| Effect of transplanting date on dry matter, yield and accumulation of photothermal units in rice (*Oryza sativa*) at Nagpur | S. R. Ghadekar, D. K. Morey and Y. P. Zilpe | 339 |
| Yield attributes of wheat (*Triticum aestivum* and *T. durum*) varieties under different sowing dates | A. D. Saini, V. K. Dadhwal and R. Nanda | 342 |
| Performance of different forage legume–*Rhizobium* symbiotic systems under saline condition | Lalita Batra and S. K. Ghai | 350 |
| Chemical behaviour of some salt-affected soils in Malaprabha and Ghataprabha river-valley projects | S. C. Kotur and T. Seshagiri Rao | 354 |
| Response of groundnut (*Arachis hypogaea*) varieties to phosphorus fertilization | Y. S. Chauhan, V. K. Jain, M. P. Khandekar and P. C. Jain | 359 |
| Effect of organic manures on economy and utilization of nitrogen by potato (*Solanum tuberosum*) and succeeding wheat (*Triticum aestivum*) | R. C. Sharma, P. M. Govindakrishnan, R. P. Singh and H. C. Sharma | 362 |
| Weed management in roselle (*Hibiscus sabdariffa* var. *altissima*) grown in different soil types | A. K. Bhattacharjee, Nirmalendu Mukerjee and V. N. Saraswat | 368 |
| A climatic approach for forecasting *Myzus persicae* on crop of seed potato (*Solanum tuberosum*) at Hisar | R. K. Kashyap and O. P. Bishnoi | 374 |
| Chemical control of apple-fruit moth, *Argyresthia conjugella* (Lepidoptera : Yponomeutidae) | J. P. Sharma, D. R. Khajuria and G. S. Dogra | 378 |
| Performance of some jujube (*Ziziphus mauritiana*) cultivars in semi-arid zone of Punjab | Room Singh and N. S. Tomer | 382 |
| Effect of nitrogen on yield of *Solanum laciniatum* | S. D. Bhardwaj and A. K. Sharma | 384 |
| Yield stability of different genotypes of rice (*Oryza sativa*) as influenced by season | I. K. Jaggi, R. O. Das and D. C. Bisen | 389 |

| | | PAGE |
|---|---|---|
| Path analysis in segregating generation of sunflower (*Helianthus annuus*) | S. K. Naskar, P. K. Bhowmik and K. Bhadra | 393 |
| Variability in castor (*Ricinus communis*) | P. S. Patel and S. N. Jaimini | 394 |
| Analysis of generation means for metric traits in upland cotton (*Gossypium hirsutum*) | H. S. Kalsy and H. R. Garg | 397 |
| Effect of disruptive selection on seed-oil content in upland cotton (*Gossypium hirsutum*) | S. S. Narayanan, Phundan Singh and V. V. Singh | 399 |
| Variability and correlations for biomass attributes in upland cotton (*Gossypium hirsutum*) | V. V. Singh | 401 |
| Genetics of yield and its components in eggplant (*Solanum melongena*) | N. D. Singh and R. K. Mital | 402 |
| Effect of sugary secretion on winter sorghum (*Sorghum bicolor*) with different N levels in drylands | C. B. Patil and K. B. Chavan | 404 |
| Prediction of yield potential of potato (*Solanum tuberosum*) through leaf and petiole potassium | U. C. Sharma and B. R. Arora | 406 |
| Effect of row spacing on forage yield of *Chrysopogon fulvus* | G. K. Dwivedi, K. C. Kanodia and R. P. Singh | 408 |
| Effect of cobalt nutrition on yield of pigeonpea (*Cajanus cajan*) | A. Rajender and A. Shiv Raj | 410 |
| Allelopathic effects of leachates and extracts of *Pinus roxburghii* on four legumes in Kumaun Himalayas | S. K. Singh and K. R. Verma | 412 |
| Incidence of *Atherigona miliaceae* (Diptera : Anthomyiidae) on proso millet (*Panicum miliaceum*) in Rajasthan | S. K. Verma | 414 |
| Relationship between meteorological parameters and occurrence of *Nephotettix* species | S. K. Shrivastava and K. C. Mathur | 416 |
| Residues of carbaryl and lindane in pointed gourd (*Trichosanthes dioica*) and its soil | R. C. Jhala, A. H. Shah and Z. P. Patel | 418 |
| Effect of saline water on performance of saplings of jujube (*Ziziphus mauritiana*), Indian cherry (*Cordia dichotoma* var. *wallichii*) and pomegranate (*Punica granatum*) at nursery stage | B. L. Jain and H. C. Dass | 420 |

| | | PAGE |
|---|---|---|
| Yield improvement through simultaneous improvement of yield components in upland cotton (*Gossypium hirsutum*) | Munshi Singh, S. N. Bhardwaj, A. D. Saini and V. P. Singh | 425 |
| Associative effect of intercropped legumes on pearlmillet (*Pennisetum glaucum*) and their residual effect on nitrogen nutrition of succeeding bread wheat (*Triticum aestivum*) | B. P. Patil and Mahendra Pal | 429 |
| Production potential of legumes, oilseeds and cereal in intercropping system with pigeonpea (*Cajanus cajan*) | A. Rafey and U. K. Verma | 433 |
| Fenugreek (*Trigonella foenum-graecum*) response to sowing date and spacing | D. S. Bhati | 437 |
| Efficacy of herbicides for weed control in winter French bean (*Phaseolus vulgaris*) | Masood Ali | 440 |
| Effect of irrigation and nitrogen fertilization on yield, quality, N uptake and water use of watermelon (*Citrullus lanatus*) | D. M. Hegde | 444 |
| Effect of dose and frequency of gypsum application on properties of sodic soil and on performance of rice (*Oryza sativa*) and bread wheat (*Triticum aestivum*) | Ram K. Gupta, O. P. Sharma and S. K. Dubey | 449 |
| Effect of residual sodium carbonate in irrigation water on soil and bread wheat (*Triticum aestivum*) | R. P. S. Chauhan, Bhudayal and C. P. S. Chauhan | 454 |
| A method to overcome the problem of spraying pigeonpea (*Cajanus cajan*) | R. Chauhan, Ombir, V. P. Singh and Y. S. Tomer | 459 |
| Chemical control of *Pseudoperonospora cubensis* causing downy mildew on sponge gourd (*Luffa cylindrica*) | L. S. Suhag, S. K. Maheshwari and J. C. Duhan | 462 |
| Pre-harvest forecast models for prediction of sugarcane yield | B. H. Singh and S. R. Bapat | 465 |
| Genetic relationship among yield components in rice (*Oryza sativa*) | G. S. V. Prasad, A. S. R. Prasad, M. V. S. Sastry and T. E. Srinivasan | 470 |
| Genotype × environment interaction in rice (*Oryza sativa*) | Narendra Kulkarni, P. Pratap Reddy, D. V. S. Raghavender Rao and G. Madhusudan Rao | 473 |

| | | PAGE |
|---|---|---|
| Combining ability for yield and its contributing characters in macaroni wheat (*Triticum durum*) | S. R. Maloo | 475 |
| Gene effects and heterosis in castor (*Ricinus communis*) | H. C. Pathak, S. K. Dixit and P. G. Patel | 476 |
| Combining ability in pea (*Pisum sativum*) | P. K. Moitra, S. P. Singh and A. K. Mehta | 479 |
| Phenotypic stability in onion (*Allium cepa*) | A. S. Sidhu, M. L. Chadha, Surjan Singh and M. R. Thakur | 481 |
| Genetic variability and path analysis in white popinac (*Leucaena latisiliqua*) | V. K. Gupta aad B. D. Patil | 483 |
| Influence of irrigation and nitrogen on water use and growth of Kinnow mandarin (*Citrus reticulata*) | K. J. A. Mageed, B. B. Sharma and A. K. Sinha | 484 |
| Effect of fertilizers on growth, root and alkaloid yield of rauvolfia (*Rauvolfia serpentina*) | S. K. Maheshwari, S. Yadav, S. K. Gangrade and K. C. Trivedi | 487 |
| Physiological analysis of growth and yield in onion (*Allium cepa*) | N. K. Srinivasa Rao | 489 |
| Consistency analysis of important chemical constituents of various tomato (*Lycopersicum esculentum*) varieties | K. L. Bajaj, Gurdeep Kaur, Surjan Singh and K. S. Nandpuri | 942 |
| Weather factors influencing outbreak of yellow rice-borer (*Scirpophaga incertulas*) on rainy-season rice (*Oryza sativa*) | P. S. Prakasa Rao and G. Padhi | 494 |
| Efficacy of systemic insecticides against cotton jassid (*Amrasca devastans*) and their effect on non-target organisms in upland cotton (*Gossypium hirsutum*) | B. Sarath Babu and G. P. Gupta | 496 |
| Effect of some insecticides on population of leafhoppers *Amrasca devastans* and *Cestius phycitis* infesting eggplant (*Solanum melongena*) | T. K. Banerjee and D. Raychaudhuri | 500 |
| Preliminary trial on chemical control of mango hoppers (*Amritodus atkinsoni* and *Idioscopus niveosparsus*) in laboratory | R. D. Pingale and S. P. Patil | 502 |

| | | PAGE |
|---|---|---|
| Build up of hexachlorocyclohexane residue in rice (*Oryza sativa*), rice-field soil and water from different formulations | P. Sukul, S. Pal, A. Chakravarty, A. Bhattacharyya, A. K. Das, A. Chowdhury and N. Adityachaudhury | 504 |
| Residues of pirimiphos-methyl and malathion spray on greengram (*Vigna radiata*) stored in jute sacks | A. K. Dikshit | 506 |
| Effect of long-term application of fertilizers, and organic and inorganic amendments under continuous cropping on soil physical and chemical properties in an Alfisol | C. L. Acharya, S. K. Bishnoi and H. S. Yaduvanshi | 509 |
| Response of rainfed pearlmillet (*Pennisetum glaucum*) to plant density and moisture conservation | S. K. Pandey, S. K. Kaushik and R. C. Gautam | 517 |
| Response of autumn potato (*Solanum tuberosum*) to irrigation and straw mulching in northern India | Baldev Singh, T. S. Aujla, B. S. Sandhu and K. L. Khera | 521 |
| Evaluation of sorghum (*Sorghum bicolor*) and safflower (*Carthamus tinctorius*) sequence cropping on dryland | S. M. Nikam, A. V. Tendulkar and A. B. Deokar | 525 |
| Effect of parasitism by *Cotesia marginiventris* (Cresson) on consumption and utilization of artificial diet by larvae of *Spodoptera litura* (Lepidoptera : Noctuidae) | S. K. Jalali, S. P. Singh and Chandish R. Ballal | 529 |
| Insecticidal control of bollworms on cotton (*Gossypium hirsutum*) | D. Thangaraju, S. Parameswaran and S. Jayaraj | 533 |
| Screening of rice (*Oryza sativa*) germplasm for resistance to sheath blight (*Rhizoctonia solani*) | B. K. Borthakur and S. K. Addy | 537 |
| Resistance of 'BJ 1' rice (*Oryza sativa*) to bacterial blight (*Xanthomonas campestris* pv *oryzae*) and genetic analysis of brown planthopper (*Nilaparvata lugens*) resistance | R. K. Sahu | 539 |
| Metroglyph analysis of yield attributes in certain tall *indica* rice (*Oryza sativa*) varieties in north Kerala | N. K. Vijayakumar and N. N. Ramankutty | 542 |
| Effect of moisture stress on yield performance in barley (*Hordeum vulgare*) | R. K. Rana, S. C. Sharma, R. S. Malik and Naresh Kumar | 543 |

| | | PAGE |
|---|---|---|
| Reaction of pigeonpea (*Cajanus cajan*) lines to wilt pathogen, *Fusarium udum* | A. Das Gupta and P. K. Sen Gupta | 546 |
| Resistance of linseed (*Linum usitatissimum*) germplasm to rust (*Melampsora lini*) and powdery mildew (*Oidium lini*) | Rajendra Prasad, M. Rai and S. A. Kerkhi | 548 |
| Genetics of resistance to *Alternaria lini* in linseed (*Linum usitatissimum*) | N. K. Singh and Y. S. Chauhan | 550 |
| Heritability and genetic advance in linseed (*Linum usitatissimum*) | Narendra Singh and N. N. Dikshit | 552 |
| Variability of oil and lint traits and effects of harvest in upland cotton (*Gossypium hirsutum*) | P. D. Kashalkar, D. G. Weginwar and R. G. Dani | 554 |
| Association of fibre yield with reproducing parts in ramie (*Boehmeria nivea*) | S. L. Ahuja | 556 |
| Effect of temperature and photosynthetically active radiation on boll weight in upland cotton (*Gossypium hirsutum*) | S. N. Bhardwaj, A. D. Saini, Munshi Singh and K. P. Singh | 557 |
| Physico-chemical characters of cigar-wrapper tobacco (*Nicotiana tabacum*) grades | V. K. Chaturvedi | 559 |
| Comparison of kinetic parameters of batch and continuous ethanol production from juice of sugarbeet (*Beta vulgaris*) | Lata R. Nain and R. S. Rana | 560 |
| Tillage requirements for seed production of potato (*Solanum tuberosum*) | V. S. Kushwah and S. S. Saini | 563 |
| Response of some vegetable crops to different sources and methods of application of zinc | B. R. V. Iyengar and M. Edward Raja | 565 |
| Effect of nitrogen application and row spacing on coriander (*Coriandrum sativum*) production under irrigated condition in semi-arid Rajasthan | D. S. Bhati | 568 |
| Determination of representative leaves for measuring leaf area in cashew (*Anacardium occidentale*) germplasm | S. Bhagavan and K. V. J. Mohan | 570 |
| Relationship between soil properties and leaf nutrient of 'Cleopatra' mandarin (*Citrus reticulata*) seedlings | T. O. Oseni | 571 |

| | | PAGE |
|---|---|---|
| Index to quantify black-tip disorder of mango (*Mangifera indica*) in orchards and its relationship with brick kilns | S. E. S. A. Khader, M. S. Rajput and P. P. Biswas | 573 |
| Screening of peach (*Prunus persica*) cultivars for their reaction against leaf curl (*Taphrina deformans*) | M. C. Nautiyal, S. Prakash and A. Kumar | 575 |
| Effect of sodicity on seed germination and seedling growth of bael (*Aegle marmelos*) | G. Pandey and R. K. Pathak | 577 |
| Antifeedant activity of oil fraction of seed of some leguminous plants against *Diacrisia obliqua* | Kishor Kumar Mohanty, D. P. Chakraborty and Shyamali Roy | 579 |
| Effect of constant temperature and relative humidity on the mortality of diapausing larvae of pink bollworm (*Pectinophora gossypiella*) of upland cotton (*Gossypium hirsutum*) | G. S. Simwat, Ashok K. Dhawan and A. S. Sidhu | 580 |
| Effect of date of sowing on the productivity of Indian mustard (*Brassica juncea* subsp. *juncea*) in the short and mild winter of the Gangetic plains of West Bengal | R. K. Ghosh and B. N. Chatterjee | 589 |
| Effect of pre-sowing seed treatment on the yield of grasspea (*Lathyrus sativus*) sown as *utera* crop in field of rice (*Oryza sativa*) | K. K. Bhattacharyya, Bikash K. Mondal, A. M. Paul, B. J. Sarkar and K. Sengupta | 596 |
| Response of potato (*Solanum tuberosum*) cultivars to phosphorus in acidic soils of Meghalaya | Kamla Singh and U. C. Sharma | 600 |
| Evaluation of various extractants for available sulphur in wetland rice (*Oryza sativa*) soils of Bangladesh | M. M. Islam and N. I. Bhuiyan | 603 |
| Differential sodicity tolerance of growth and yield parameters in genotypes of rice (*Oryza sativa*) | Ali Qadar | 607 |
| Early-generation selection for fruit number and fruit yield in six crosses of eggplant (*Solanum melongena*) | Narendra Kumar and Hari Har Ram | 611 |
| Reaction of pearlmillet (*Pennisetum glaucum*) varieties to khapra beetle (*Trogoderma granarium*) | S. V. Dhamdhere, R. S. Tomar, H. S. Narwaria, M. L. Sharma, M. P. Khare, U. S. Misra and U. S. Dwivedi | 615 |

| | | PAGE |
|---|---|---|
| Effect of host plants on some biological aspects of spotted pod-borer (*Maruca testulalis*) (Lepidoptera : Pyralidae) | G. V. Ramasubramanian and P. C. Sundara Babu | 618 |
| Toxicity of insecticides on eggs, nymphs and adults of tur-pod bug (*Clavigralla gibbosa*) and its ovipositional behaviour on pigeonpea (*Cajanus cajan*) | K. J. Singh, R. C. Thakur and O. P. Singh | 621 |
| Effect of common pesticides used in Nigeria on soil microbial population | O. Odeyemi, A. Salami and Esther O. Ugoji | 624 |
| Genetic variability and correlation coefficients in early rice (*Oryza sativa*) | T. Sundaram, W. Wilfred Manuel and S. Palanisamy | 629 |
| Evaluation of new male-sterile lines of sorghum (*Sorghum bicolor*) for yield and disease reaction | H. Shivanna and K. H. Anahosur | 631 |
| Genetic analysis for stem sugar in sorghum (*Sorghum bicolor*) | B. Selvi and S. Palanisamy | 632 |
| Genotype × environment interaction for seed yield in brown sarson (*Brassica rapa* var. *ulti*) | A. Henry and H. S. Daulay | 634 |
| Association and index selection for germinability by propagation methods in ramie (*Boehmeria nivea*) | S. L. Ahuja and O. P. Tuteja | 636 |
| Effect of water stress in Ethiopian mustard (*Brassica carinata*) and Indian mustard (*B. juncea* subsp. *juncea*) on infestation by *Lipaphis erysimi* | D. R. C. Bakhetia and K. S. Brar | 638 |
| Optimum use of limited irrigation water for winter crops in south-west Punjab | N. T. Singh, A. S. Sidhu, G. S. Dhaliwal and G. C. Aggarwal | 641 |
| *Rhizobium-Sclerotium* interaction in groundnut (*Arachis hypogaea*) rhizosphere | P. Bhattacharyya and N. Mukherjee | 643 |
| Biochemical changes during growth and development of linseed (*Linum usitatissimum*) | S. Dua and K. Rani | 645 |
| Fungicidal control of alternaria leaf spot (*Alternaria alternata*) of sesame (*Sesamum indicum*) | M. Achuta Rama Rao, S. Ramapandu and Y. Satyanarayana | 648 |

| | | PAGE |
|---|---|---|
| Incidence, damage and carry-over of cutworm (*Agrotis ypsilon*) attacking potato (*Solanum tuberosum*) crop in Bihar | B. B. Das and Gulab Ram | 650 |
| Biology of six syrphid predators of cabbage aphid (*Brevicoryne brassicae*) on seed crop of cauliflower (*Brassica oleracea* var. *botrytis*) | K. C. Sharma and O. P. Bhalla | 652 |
| Effect of summer application of insecticides on San Jose scale (*Quadraspidiotus perniciosus*) in orchards of apple (*Malus pumila*) | S. P. Bhardwaj | 655 |
| Effect of soil-moisture level on population of *Tylenchulus semipenetrans* infecting sour orange (*Citrus aurantium*) | M. E. Sweelam and M. R. Abo El-Ghar | 657 |
| Ovipositional host status of some weeds, wild rice (*Oryza* species) and cultivated crop plants for *Nilaparvata lugens* | S. M. Zaheruddeen and P. S. Prakasa Rao | 659 |
| Transmission of nuclear polyhedrosis virus of *Heliothis armigera* (Lepidoptera : Noctuidae) through insectivorous birds | R. V. Vyas, B. M. Parasharya and D. N. Yadav | 661 |
| Zero-energy cool chamber for maintaining post-harvest quality of carrot (*Daucus carota* var. *sativa*) | R. K. Pal and Susanta K. Roy | 665 |
| Irrigation and nitrogen requirement of bell pepper (*Capsicum annuum*) | D. M. Hegde | 668 |
| Spatial arrangement and methods of planting in cabbage (*Brassica oleracea* convar. *capitata* var. *capitata*) intercropping | R. P. Sharma, R. R. Patil and P. N. Arora | 673 |
| Effect of sowing date on heat-unit requirement and yield of winter groundnut (*Arachis hypogaea*) | S. R. Ghadekar | 678 |
| Response of wheat (*Triticum aestivum*) to boron application | A. B. Mandal and A. D. Das | 681 |
| Genotype × environment interaction for seed yield in horsegram (*Macrotyloma uniflorum*) | A. Henry and H. S. Daulay | 684 |

| | | PAGE |
|---|---|---|
| Heterosis for yield and its components, dehydration qualities and storage in onion (*Allium cepa*) | Netra Pal, Narendra Singh and B. Choudhury | 687 |
| Control of black-scurf (*Rhizoctonia solani*) and common scab (*Streptomyces scabies*) of potato (*Solanum tuberosum*) with boric acid | A. K. Somani | 693 |
| Effect of surface crust breaking on growth and yield of bread wheat (*Triticum aestivum*) in alkali soils | D. P. Sharma | 699 |
| Effect of water stress on yield attributes and yield of groundnut (*Arachis hypogaea*) | M. S. Patel and B. A. Golakiya | 701 |
| Effect of potassium fertilization on winter hardiness and yield of bread wheat (*Triticum aestivum*) and its residual effect on greengram (*Vigna radiata*) under rainfed conditions | G. H. Nakashgir, G. M. Khan, M. M. Rafique and S. A. Wani | 704 |
| Response of vegetable french bean (*Phaseolus vulgaris*) to nitrogen and phosphorus fertilization | K. Srinivas and L. B. Naik | 707 |
| Effect of various cropping patterns including cereals, pulses and oilseeds on chemical properties of soil | H. C. Thakur and N. N. Sharma | 708 |
| Relationship between root weight and plant characters in rice (*Oryza sativa*) in different seasons | M. Abdul Salam and S. Subramanian | 710 |
| Evaluation of grain protein content in selected high-protein genotypes of barley (*Hordeum vulgare*) grown under low and high nitrogen fertility | Raghuveer Polisetty, R. Chandra, L. V. S. Sastry, B. N. Murthy and G. S. Sirohi | 711 |
| Morphology of spike in relation to resistance to loose smut (*Ustilago tritici*) in wheat (*Triticum aestivum*) | D. K. Pandey and P. L. Gautam | 713 |
| Heterosis and combining ability in relation to cytoplasmic diversity in sorghum (*Sorghum bicolor*) | A. Ganga Kishan and S. T. Borikar | 715 |
| Combining ability in grain sorghum (*Sorghum bicolor*) | R. H. Patel, K. B. Desai, M. S. Desai and N. D. Modi | 717 |
| Genetic variability in tomato (*Lycopersicum esculentum*) | P. K. Singh, R. K. Singh, B. C. Saha and Rajesh Kumar | 718 |

| | | PAGE |
|---|---|---|
| Resistance of chilli (*Capsicum* species) to tobacco mosaic and cucumber mosaic viruses | N. L. Peshney and P. G. Moghe | 720 |
| Effectiveness of selection at seedling and single-row stages in cassava (*Manihot esculenta*) | P. G. Rajendran, K. R. Lakshmi and G. G. Nayar | 722 |
| Evaluation of exotic cultivars for yield and chemical characters of globe artichoke (*Cynara scolymus*) in Kashmir valley | P. N. Pandita, R. K. Ogra and Archana Kaul | 724 |
| Estimation of leaf area of french bean (*Phaseolus vulgaris*) using linear measurements | Ajambar Rai, Percival B. Alipit and Marilyn B. Toledo | 727 |
| Role of nitrogen, phosphorus and potassium fertilizers and organic amendments in cumin (*Cuminum cyminum*) wilt incited by *Fusarium oxysporum* f. sp. *cumini* | R. S. Champawat and V. N. Pathak | 728 |
| Pests of milk tree (*Manilkara hexandra*), a rootstock for sapodilla (*Manilkara achras*) | R. C. Jhala, Z. P. Patel and A. H. Shah | 730 |
| Effect of fertilizers on incidence of shoot-and-fruit borer on okra (*Abelmoschus esculentus*) | K. K. Kumar and K. C. Devaraj Urs | 732 |
| Effective insecticidal schedule for control of major insect pests of rice (*Oryza sativa*) | N. P. V. Vijaya Kumar, M. M. Krishna Murthy and G. P. V. Reddy | 734 |
| Determination of waiting period of phosphamidon in okra (*Abelmoschus esculentus*) | P. K. Vaidya and Vikram Singh | 736 |
| New sources of early-maturing germplasm in pearlmillet (*Pennisetum glaucum*) | S. Appa Rao, Melak H. Mengesha and C. Rajagopal Reddy | 743 |
| Heterosis in intervarietal crosses of bell pepper (*Capsicum annuum* var. *grossum*) and hot chilli (*C. annuum* var. *fasciculatum*) | Pious Thomas and K. V. Peter | 747 |
| Effect of drainage and sowing methods on yield and its attributes of maize (*Zea mays*) | U. K. Prasad, R. D. Pandey and N. N. Sharma | 751 |
| Response of Indian mustard (*Brassica juncea* subsp. *juncea*) to sulphur fertilization | V. K. Singh, Vinay Singh and Mahabir Singh | 754 |

| | | PAGE |
|---|---|---|
| Effect of phosphorus in presence and absence of sulphur and magnesium on yield and uptake of P, S and Mg by 'GAUG 10' groundnut (*Arachis hypogaea*) | G. L. Maliwal and N. K. Tank | 757 |
| Forms of phosphorus in soils of Karnataka | V. S. Doddamani and T. Seshagiri Rao | 760 |
| Reprocessing of infected parboiled rice (*Oryza sativa*) | K. Singaravadivel and S. Anthoni Raj | 766 |
| Effect of water stress on grain growth of glossy and non-glossy varieties of grain sorghum (*Sorghum bicolor*) | C. L. Narasimha Rao and A. Shivraj | 770 |
| Yield and quality of summer groundnut (*Arachis hypogaea*) as influenced by growth retardants and promoters | M. S. Patel, J. D. Gundalia, R. L. Davaria and P. G. Vadher | 773 |
| Performance of *Rhizobium japonicum* strains tolerant to high temperature and low soil moisture | V. R. Balasundaram | 776 |
| Optimum compost depth in polythene bags for *Agaricus bisporus* cultivation | R. C. Upadhyay and B. Vijay | 778 |
| Incidence of maize stalk-borer (*Chilo partellus*) in relation to fertilizer level | R. N. Chaudhary and V. K. Sharma | 780 |
| Efficacy of synthetic pyrethroids against rice hispa (*Dicladispa armigera*) in field | N. V. Krishnaiah, Mangal Sain and I. C. Pasalu | 781 |
| Efficacy of new insecticides for control of insect pests of spring okra (*Abelmoschus esculentus*) | Gajendra Singh and P. N. Misra | 783 |
| Food utilization by *Heliothis armigera* (Lepidoptera : Noctuidae) | P. R. Srinivas | 785 |
| Dissipation of fenitrothion and monocrotophos on pods of pigeonpea (*Cajanus cajan*) | Pashupati Singh, S. F. Hameed and I. P. Singh | 787 |
| Diallel-cross analysis for yield and its components in bread wheat (*Triticum aestivum*) | K. M. S. Raghuvanshi, S. P. Singh, S. K. Rao and C. B. Singh | 789 |
| Combining-ability analysis for protein and *in-vitro* dry-matter digestibility in forage sorghum (*Sorghum bicolor*) | R. S. Khatri and D. S. Jatasra | 792 |

| | | PAGE |
|---|---|---|
| Genotype × environment interaction for seed yield in Indian mustard (*Brassica juncea* subsp. *juncea*) | A. Henry and H. S. Daulay | 794 |
| Heritability and genetic advance for yield and its components in diverse crosses of blackgram (*Vigna mungo*) | R. N. Sharma and S. K. Rao | 795 |
| Genetic variability of yield and its components in horsegram (*Macrotyloma uniflorum*) | D. Suraiya, S. T. Mercy, D. Chandramony and N. K. Nayar | 798 |
| Correlation and path analysis in eggplant (*Solanum melongena*) | S. C. Khurana, G. Kalloo, C. B. Singh and K. K. Thakral | 799 |
| Association in parents and segregating populations of pea (*Pisum sativum*) | S. P. Singh and Ajit Lavania | 801 |
| Stability of cotton (*Gossypium* species) in relation to soil depth under rainfed condition | N. N. Nimbole | 803 |
| Transfer of cytoplasmic male sterility in upland cotton (*Gossypium hirsutum*) | Sunil Seth, D. K. Jain and M. S. Kairon | 805 |
| Metroglyph and index-score character analysis of some exotic sugarcane hybrids | Sardul Singh Gill and B. K. Tripathi | 806 |
| Energetics of wheat (*Triticum aestivum*) production as influenced by tillage and nitrogen management | U. N. Verma, V. C. Srivastava and U. K. Verma | 813 |
| Performance and quality of summer cowpea (*Vigna unguiculata* subsp. *cylindrica*), soybean (*Glycine max*) and pearlmillet (*Pennisetum glaucum*) grown in mixture and intercropping under different moisture regimes | S. S. Narwal, S. Singh, D. S. Malik and P. C. Gupta | 817 |
| Response of rice (*Oryza sativa*) and wheat (*Triticum aestivum*) to phosphorus and zinc fertilization in sodic soil | M. V. Singh | 823 |
| Timing of scheduled insecticide application for control of leafhopper (*Amrasca biguttula biguttula*) on okra (*Abelmoschus esculentus*) | K. Srinivasan and N. K. Krishnakumar | 828 |

| | | PAGE |
|---|---|---|
| Effect of adjoining host crops, date of last cut and natural enemies on *Heliothis armigera* infesting seed crop of Egyptian clover (*Trifolium alexandrinum*) | J. S. Dhaliwal and D. S. Sidhu | 832 |
| Improvement for resistance to maydis leaf-blight (*Drechslera maydis*) and maize-borer (*Chilo partellus*) in maize (*Zea mays*) | S. K. Dey, B. S. Dhillon, V. K. Saxena, Uma Kanta and A. S. Khehra | 837 |
| Inheritance and linkage relationships of qualitative characters in pearlmillet (*Pennisetum glaucum*) | S. Appa Rao, Melak H. Mengesha and C. Rajagopal Reddy | 840 |
| Field screening of mothbean (*Vigna aconitifolia*) for susceptibility to insect pests and diseases | R. K. Dabi and H. N. Gour | 843 |
| Economic potential of wheat (*Triticum aestivum*)–pearlmillet (*Pennisetum glaucum*) rotation as affected by soil amendments and phosphate fertilization under saline-sodic conditions | G. L. Keshwa and G. D. Singh | 845 |
| Economy of fertilizer nitrogen through green-manuring in rice (*Oryza sativa*) | D. S. Rana, Hargopal Singh, K. N. Sharma and A. L. Bhandari | 848 |
| Utilization of modified urea in wetland rice (*Oryza sativa*) | S. S. Tomar and J. K. Verma | 849 |
| Response of rice (*Oryza sativa*) to potassium application under two water-management practices | B. P. Singh and B. N. Singh | 851 |
| Effect of plant density, and level and proportion of nitrogen fertilization on growth, yield and yield components of winter maize (*Zea mays*) | A. S. Bangarwa, M. S. Kairon and K. P. Singh | 854 |
| Critical concentration of zinc for soybean (*Glycine max*) grown in calcareous soil of north Indian plains | A. N. Jha and A. S. Chandel | 857 |
| Effect of time of weed removal on yield of greengram (*Vigna radiata*) | Shiv Kumar and M. S. Kairon | 859 |
| Effect of nitrogen and phosphorous on fruit yield and quality of sponge gourd (*Luffa aegyptiaca*) | Satish Siyag and S. K. Arora | 860 |

| | | PAGE |
|---|---|---|
| Estimation of leaf area of two okra (*Abelmoschus esculentus*) varieties through leaf characteristics | S. N. Asoegwu | 862 |
| Chemical control of pod-borer complex of greengram (*Vigna radiata*) | N. S. Bhat, M. Manjunatha, G. T. T. Raju, G. G. Nagabhushana and V. P. Deshpande | 864 |
| Seasonal incidence of shoot-and-fruit borer (*Leucinodes orbonalis*) on egg-plant (*Solanum melongena*) in Gujarat | K. V. Thanki and J. R. Patel | 867 |
| Aldicarb residues in soil and their translocation into potato (*Solanum tuberosum*) tubers | T. S. Kathpal, G. Singh, G. S. Yadav, R. S. Jaglan and Y. S. Malik | 868 |
| Host range of solitary *Platygaster* species, a parasitoid of rice gall-midge (*Orseolia oryzae*) | K. C. Mathur, P. K. Das and S. Sasmal | 871 |
| Effect of plant species on post-embryonic development and survival of armyworm (*Mythimna separata*) | Ram Singh and J. P. Chaudhary | 873 |
| Effect of refrigeration of puparia on adult emergence and pupal period of *Eucelatoria bryani* | Shekharappa, K. Jairao and Yelshetty Suhas | 875 |
| Estimation of genetic parameters in grain sorghum (*Sorghum bicolor*) | R. H. Patel, K. B. Desai, M. S. Desai and K. G. Patel | 877 |
| Involvement of ethylene with new wilt of cotton (*Gossypium* species) | R. K. Sahay, N. L. Bhale, R. Pundarikakshudu and M. R. K. Rao | 885 |
| Criteria for evaluating resistance of grain sorghum (*Sorghum bicolor*) to striga (*Striga hermonthica*) | N. A. Nafie and A. E. Abdalla | 888 |
| Screening for resistance to sterility mosaic disease of pigeonpea (*Cajanus cajan*) | R. P. Gupta, B. B. Singh, D. P. Singh aud Vinod Singh | 893 |
| Economics of rice (*Oryza sativa*) cultivation in different seasons as influenced by levels of nitrogen, zinc sulphate and granular insecticides | M. Abdul Salam and S. Subramanian | 895 |
| Effect of gamma irradiation, evaporation retardants and transpiration suppressants on grain yield, nutrient uptake and moisture-use efficiency on bread wheat (*Triticum aestivum*) | D. K. Dash, Avijit Sen and N. M. Misra | 901 |

| | | PAGE |
|---|---|---|
| Efficiency of biomass production in intercropped pearlmillet (*Pennisetum glaucum*)–bread wheat (*Triticum aestivum*)–greengram (*Vigna radiata*) cropping system | Mahendra Pal and B. P. Patil | 905 |
| Effect of irrigation scheduling and phosphorus doses on chickpea (*Cicer arietinum*) in deep Vertisols | P. S. Kulhare, K. P. Tiwari, R. N. Saran and J. P. Dixit | 911 |
| Response of ratoon crop of sugarcane to gap filling and fertilization | U. P. Sinha, K. C. Jha, S. Zama and H. Singh | 915 |
| Oviposition and larval behaviour of stalk borer *Chilo partellus* on susceptible and resistant varieties of maize (*Zea mays*) | Harish Kumar | 918 |
| Differential susceptibility of potato (*Solanum tuberosum*) cultivars to zinc stress and nutrient interaction | S. P. Trehan, U. C. Sharma and J. S. Grewal | 922 |
| Evaluation of passive evaporative cooling for storage of potato (*Solanum tuberosum*) | H. N. Kaul and Ashiv Mehta | 928 |
| Effect of cutting angle of shovel on unit draft and penetration | A. C. Varshney and K. V. Patel | 932 |
| Leaf-disc, a new non-destructive method of sampling mango leaf | K. Malla Reddy and A. Rameshwar | 935 |
| Effect of benzyladenine on endosperm-free lipid pool and its fractions in grains of bread wheat (*Triticum aestivum*) | P. C. Pande and S. N. Bhardwaj | 936 |
| Phosphate uptake by wheat (*Triticum* species) varieties grown under irrigated and rainfed regions of Maharashtra | A. S. Nirale | 939 |
| Rate and duration of pod-filling and seed filling periods of greengram (*Vigna radiata*) | R. Nanda and A. D. Saini | 941 |
| Effect of levels of nitrogen and phosphorus on forage yield of rainfed 'MP Chari' sorghum (*Sorghum bicolor*) | M. H. Shah, A. S. Bali, K. N. Singh and P. K. Koul | 945 |
| Effect of nitrogen, phosphorus and zinc on yield of cauliflower (*Brassica oleracea* convar. *botrytis* var. *botrytis*) | Dharam Veer Balyan and B. S. Dhankhar | 947 |

| | | PAGE |
|---|---|---|
| Performance of arum (*Colocasia esculenta*) under selected cultural management practices | M. Prabhakar and G. M. Nair | 949 |
| Management of fruit diseases of guava (*Psidium guajava*) through fungicidal sprays | R. D. Rawal and B. A. Ullasa | 950 |
| Phytotoxicity of carbaryl to soybean (*Glycine max*) and pigeonpea (*Cajanus cajan*) | A. K. Awasthi and S. M. Vaishampayan | 953 |
| Inheritance of shootfly (*Atherigona soccata*) resistance in $F_2$ diallel crosses of sorghum (*Sorghum bicolor*) | V. S. Nimbalkar and D. R. Bapat | 955 |
| Metroglyph analysis in persian clover (*Trifolium resupinatum*) | R. K. Gumber and M. S. Sohoo | 958 |
| Book Reviews | | 83, 157, 245, 333, 422, 583, 739, 808, 880, 960 |